

APPLICATION FORM (Deadline March 1st , 2021)

SHORT CV

Camille Vrijghem

Master in Dentistry (2013- 2018) – KU Leuven

Master after master in Periodontology and Oral Implantology (2018-2021) – Ghent University

Publication:

Vrijghem C, Cosyn J, Franssen J, Christiaens V. A questionnaire-based crossover study on the association of microbiological testing with prescription of systemic antibiotics following initial periodontal therapy. *J Clin Periodontol.* 2021 Mar 9. doi: 10.1111/jcpe.13453. Epub ahead of print. PMID: 33751614.

COMPOSITION OF THE TEAM (if relevant)

/

ABSTRACT

Recessions in the anterior mandibular region after previous orthodontic treatment are a well-known complication. Prevalence according to Vasconcelos and colleagues goes up to 10,3% and is predominantly present on the central incisors. When, in these cases, the buccal bone is missing, first choice of treatment is orthodontic retrusion. In this patient, a recession type 2 (Cairo 2011), there was not enough mandibular bone to retract the lower central incisors orthodontically. Therefore, the recession was covered with a coronally advanced flap combined with a connective tissue graft and EMD. (Stefanini M. & Zucchelli G.)

Stefanini M, Zucchelli G, Marzadori M, de Sanctis M. Coronally Advanced Flap with Site-Specific Application of Connective Tissue Graft for the Treatment of Multiple Adjacent Gingival Recessions: A 3-Year Follow-Up Case Series. Int J Periodontics Restorative Dent. 2018 Jan/Feb;38(1):25-33. doi: 10.11607/prd.3438. PMID: 29240201.



2 weeks post-operative



1 month post-operative

